

Predicting the solar maximum with the rising rate

Z. L. Du and H. N. Wang

*Key Laboratory of Solar Activity, National Astronomical Observatories, Chinese Academy of Sciences,
Beijing 100012, China*

zldu@nao.cas.cn

ABSTRACT

The growth rate of solar activity in the early phase of a solar cycle has been known to be well correlated with the subsequent amplitude (solar maximum). It provides very useful information for a new solar cycle as its variation reflects the temporal evolution of the dynamic process of solar magnetic activities from the initial phase to the peak phase of the cycle. The correlation coefficient between the solar maximum (β_a) and the rising rate (β_a) at Δm months after the solar minimum ($R_{\min}$) is studied and shown to increase as the cycle progresses with an inflection point ($r = 0.83$) at about $\Delta m = 20$ months. The prediction error of $R_{\max}$ based on β_a is found within estimation at the 90% level of confidence and the relative prediction error will be less than 20% when $\Delta m \geq 20$. From the above relationship, the current cycle (24) is preliminarily predicted to peak around October, 2013 with a size of $R_{\max} = 84 \pm 33$ at the 90% level of confidence.

Subject headings: solar physics; solar activity; sun spots; solar cycles

1. Introduction

The Waldmeier effect that stronger cycles tend to rise faster is a well known fact in solar activity (Waldmeier 1939; Hathaway et al. 2002; Du et al. 2009a). The growth rate of solar activity (R_z) in the early phase of a solar cycle, defined as the ratio of a given increment ($\Delta R_z = 20$) between two certain levels ($R_{z1} = 30$ and $R_{z2} = 50$) over the corresponding elapsed time (Δt), was found to be highly correlated ($r > 0.8$) with the subsequent amplitude (Cameron & Schüssler 2008). Therefore, the strength of a new cycle should be rationally predicted by the above relation. The problem is that if and at what month after the start of a new cycle the strength of the cycle can be well estimated from the early information of the cycle.

This paper studies the variation of the correlation between the maximum amplitude ($R_{\max}$) of a solar cycle and the rising rate (β_a) as a function of Δm months entering the cycle and analyzes the predictive power of β_a on $R_{\max}$ in order to find out at what month $R_{\max}$ can be well estimated by β_a . The results are shown in the following section. β_a is defined as the ratio of the increment

of R_z from the minimum ($R_{\min}$) over the elapsed time (Δm months). The temporal variation in the correlation coefficient (r) between $R_{\max}$ and β_a is analyzed in Section 2.1, showing that r is very low near the initial phase ($r < 0.5$ if $\Delta m \leq 10$) and significant only at a few months after the start of the cycle ($r > 0.8$ if $\Delta m \geq 19$). The predictive power of β_a on $R_{\max}$ as the cycle progresses is analyzed in Section 2.2, indicating that the relative prediction error of $R_{\max}$ is very small for almost all Δm in some cycles and smaller than 20% at some (about twenty) months after the start in other cycles. The peak size and its timing of cycle 24 are estimated in Section 3, followed by conclusions in Section 4.

2. Data and Analysis

The data used in the present study are the smoothed monthly mean international sunspot number (R_z)¹ from July, 1749 to February, 2011. The rising rate is defined as the ratio, $\beta_a = (R_z(\Delta m) - R_{\min})/\Delta m$, of the increment of R_z

¹<http://www.ngdc.noaa.gov/stp/spaceweather.html>

Table 1: Parameters and results

n	$R_{\min}$	$R_{\max}$	T_a	$\beta_a(27, n)$	E (E_t)	cc (cc_t)
1	8.4	86.5	76	0.74		
2	11.2	115.8	40	2.23		
3	7.2	158.5	35	3.94		
4	9.5	141.2	41	3.64		
5	3.2	49.2	82	0.45		
6	0.0	48.7	70	0.20		
7	0.1	71.5	79	0.63	0.18(0.80)	-0.15(-0.98)
8	7.3	146.9	40	3.58	0.15(0.27)	-0.90(-0.98)
9	10.6	131.9	55	1.24	0.32(0.31)	+0.39(-0.88)
10	3.2	98.0	50	1.54	0.13(0.47)	+0.11(-0.98)
11	5.2	140.3	41	2.49	0.09(0.32)	-0.32(-0.98)
12	2.2	74.6	60	1.76	0.39(0.57)	-0.35(-0.98)
13	5.0	87.9	47	2.34	0.30(0.49)	+0.18(-0.98)
14	2.7	64.2	49	1.42	0.43(0.67)	-0.65(-0.96)
15	1.5	105.4	49	1.94	0.09(0.41)	-0.01(-0.93)
16	5.6	78.1	57	1.93	0.22(0.52)	-0.30(-0.95)
17	3.5	119.2	43	1.99	0.15(0.35)	-0.29(-0.96)
18	7.7	151.8	39	2.73	0.21(0.28)	-0.81(-0.98)
19	3.4	201.3	47	5.26	0.19(0.20)	-0.96(-0.95)
20	9.6	110.6	49	2.42	0.06(0.38)	-0.22(-0.97)
21	12.2	164.5	44	3.14	0.25(0.26)	-0.93(-0.98)
22	12.3	158.5	34	4.64	0.10(0.31)	+0.65(-0.98)
23	8.0	120.8	47	2.21	0.09(0.35)	-0.82(-0.98)
24	1.7	?84.0	?59	1.17		
av.	5.9	119.1	49	2.43	0.20(0.41)	-0.31(-0.96)

from the minimum ($R_{\min}$) over the elapsed time (Δm) from the start of the cycle. The rising rate is computed for each cycle n and each Δm , denoted by $\beta_a(\Delta m, n)$. The parameters are listed in Table 1 in which $R_{\min}$ and $R_{\max}$ are the minimum and maximum amplitudes of the solar cycle, respectively; T_a is the rise time from minimum to maximum; $\beta_a(27, n)$ is the value of $\beta_a(\Delta m, n)$ at the current state $\Delta m = 27$; other parameters will be described later; and the last row indicates the relevant averages over cycles $n = 7-23$.

2.1. The variation in the correlation between $R_{\max}$ and β_a as the cycle progresses

Figure 1 illustrates the variation in the correlation coefficient (r) between $R_{\max}(n)$ and $\beta_a(\Delta m, n)$ for the cycles in which $T_a(n) \geq \Delta m$ at a given Δm (using only the data in the rising phases). One can see that r varies with the progression of the cycle (Δm). A steady increasing trend is shown in r since $\Delta m = 6$: r increases from about 0.33 at $\Delta m = 6$ to about 0.83 at




Fig. 1.— Correlation coefficient (r) between $R_{\max}$ and β_a as a function of Δm .

$\Delta m = 20$ at a high speed, and increases at a smaller speed since then, showing an inflection point at $\Delta m = 20$ months, $r(20) = 0.83$. Near the initial phase ($\Delta m \leq 10$) the correlation is not strong ($r < 0.5$). The correlation coefficient between $R_{\max}$ and β_a is high enough at $\Delta m = 19$ ($r > 0.81$) months entering the solar cycle. At the current state ($\Delta m = 27$), $r(27) = 0.88$.

2.2. The predictive power of β_a on $R_{\max}$

In order to test the predictive power of β_a on $R_{\max}$ at different Δm , we use only the data up to cycle $(n-1)$ to predict $R_{\max}$ for cycle n . For a given Δm , we calculate the linear regression equation of $R_{\max}(i)$ against $\beta_a(\Delta m, i)$ for cycles $i = 1, 2, \dots, n-1$ in the form of

$$R_{\max} = A + B\beta_a, \quad (1)$$

and the standard deviation $\sigma(\Delta m, n-1)$ used to estimate the uncertainty of the prediction of $R_{\max}(n)$. Then, substituting the value of $\beta_a(\Delta m, n)$ into this equation, the $R_{\max}$ value for cycle n can be predicted, which is denoted by $R_p(\Delta m, n)$. Figure 2 shows the results for the recent nine cycles $n = 15-23$: $R_p(\Delta m, n)$ (black solid line) together with error bars $t_r(n-1)\sigma(\Delta m, n-1)$; $R_{\max}(n)$ (black horizontal long-dashed line), the actual relative prediction error (red dotted),

$$E(\Delta m, n) = \frac{|R_p(\Delta m, n) - R_{\max}(n)|}{R_{\max}(n)}; \quad (2)$$

the estimated relative prediction error (green dashed),

$$E_t(\Delta m, n) = \frac{t_r(n-1)\sigma(\Delta m, n-1)}{R_{\max}(n)}, \quad (3)$$

where $t_r(n-1)$ is the t -value at the 90% level of confidence in a student's t -distribution for $n_f = (n-3)$ degrees of freedom; and the correlation coefficient between $R_{\max}(i)$ and $\beta_a(\Delta m, i)$ (r , blue dash-dotted line, multiplied by 100 to be indicated by the right hand scale). The numbers in the figure (cc) denote the correlation coefficients between E (E_t) and r .

Figure 2(a) illustrates the results for cycle 15. It is seen that $R_{\max}$ (black horizontal long-dashed line) is all within the error bars of R_p at the 90% level of confidence (vertical lines), $E < E_t$, and $E < 20\%$ (red horizontal long-dashed line) when $\Delta m \geq 3$, although there are some fluctuations in both E and E_t . The anti-correlation coefficient between E_t and r is very strong, $cc_t = -0.93$, implying that the higher the correlation coefficient (r) between $R_{\max}$ and β_a , the smaller the estimated relative prediction error (E_t) from the extrapolation of the relationship between $R_{\max}$ and β_a . However, this is only an estimate in theory rather than in practice. In fact, the correlation coefficient between E and r is almost zero, $cc = -0.01$, implying that the actual relative prediction error is almost uncorrelated to r . That is to say it is uncertain whether a more (less) accurate prediction corresponds to a higher (lower) correlation.

In Figure 2(d) for cycle 18, we test only the results for the rising phase $\Delta m \leq T_a (= 39)$. One can see that both E and E_t decrease as Δm increases. The anti-correlation coefficient between E_t and r is also very strong, $cc_t = -0.98$. The correlation coefficient between E and r is highly negative, $cc = -0.81$, implying that a more (less) accurate prediction can be obtained from a higher (lower) correlation in this case. In cycle 18, $R_{\max}$ is always within the error bars of R_p at the 90% level of confidence, $E < E_t$. In addition, $E < 20\%$ when $\Delta m \geq 26$.

Figure 2(g) shows the results for cycle 21: $\Delta m \leq T_a (= 44)$. The results are similar to those in Fig. 2(d): both E and E_t decrease as Δm increases; the anti-correlation coefficient between E (E_t) and r is very strong, $cc = -0.93$ ($cc_t = -0.98$). At a small Δm , E is large: $E > E_t$

if $\Delta m \leq 17$. As the cycle progresses, E becomes smaller: $E < E_t$ when $\Delta m \geq 18$; $E < 20\%$ when $\Delta m \geq 26$.

The results in other cycles are similar to those above. The main conclusions in Fig. 2 may be summarized as follows.

1. $R_{\max}$ (black horizontal long-dashed) is usually within the error bars of R_p (black solid) at the 90% level of confidence (vertical lines), apart from cycles 19 and 21 when $\Delta m \leq 16$ and $\Delta m \leq 17$, respectively;
2. the estimated relative prediction error E_t (green dashed) tends to decrease as Δm increases; E_t is highly anti-correlated with the correlation coefficient r (blue dash-dotted), $cc_t \approx -0.96$;
3. the actual relative prediction error E (red dotted) varies with some fluctuations and tends to decrease as Δm increases in some cycles (for $n = 16-19, 21, 23$);
4. there is no established relationship between E and r , such as $|cc| > 0.8$ in cycles $n = 18, 19, 21$ and 23 , while $|cc| \leq 0.3$ in cycles $n = 15, 16, 17$ and 20 , and cc is even positive in cycle 22 (0.65);
5. $E < E_t$ for all Δm in cycles $n = 15-18, 20, 22$ and 23 ; $E < E_t$ for $\Delta m \geq 17$ in cycle 19, and for $\Delta m \geq 18$ in cycle 21;
6. $E < 20\%$ (red horizontal long-dashed) since a very few months entering the cycle ($\Delta m > 3$) in cycles 15, 20 and 22; $E < 20\%$ for $\Delta m \geq 34$ in cycle 16, for $\Delta m \geq 21$ in cycle 17, for $\Delta m \geq 26$ in cycle 18, for $\Delta m \geq 17$ in cycle 19, for $\Delta m \geq 26$ in cycle 21; and for $\Delta m \geq 10$ in cycle 23,

In summary, β_a behaved very well in predicting the subsequent $R_{\max}$: (i) the actual prediction error (known only when the cycle is over) is usually within estimation since about twenty months entering the cycle, $|R_p(\Delta m, n) - R_{\max}(n)| < t_r(n-1)\sigma(\Delta m, n-1)$; (ii) the relative prediction error is usually less than 20% since about twenty months into the cycle; and (iii) E tends to decrease as the cycle progresses. In some cycles ($n = 15, 20$ and 22 , see Figures 2(a), (f) and (h)), the relative prediction error is very small ($E < 10\%$) at




Fig. 2.— Predictions (R_p , black solid line, left hand scale) of $R_{\max}$ (black horizontal long-dashed line) together with error bars ($t_t(n-1)\sigma(\Delta m, n-1)$, vertical line) for cycles 15–23 (panels (a)–(i), respectively), the actual relative prediction error (E , red dotted, right hand scale), the estimated value (E_t , green dashed), and the correlation coefficient between $R_{\max}$ and β_a (r , blue dash-dotted). The numbers in the figure (cc) denote the correlation coefficients between E (E_t) and r .

a small Δm even if the correlation coefficient is low ($r < 0.5$). Similar conclusions can also be obtained in other cycles (not shown): $E < E_t$ at about $\Delta m \geq 20$; E_t is highly anti-correlated with r (cc_t); while there is no established relationship between E and r (cc). The results of $\overline{E}$ ($\overline{E}_t$) and cc (cc_t) in cycles 7–23. The results in Table 1, and the relevant averages are indicated by the last row: $\langle \overline{E} \rangle = 0.20$ ($\langle \overline{E}_t \rangle = 0.41$) and $\langle cc \rangle = -0.31$ ($\langle cc_t \rangle = -0.96$), where $\overline{E}$ is the average over Δm in a solar cycle and $\langle \overline{E} \rangle$ represents the average over cycles 7–23. Therefore, a higher (lower) correlation coefficient does not necessarily yield a more (less) accurate prediction (Du et al. 2009b; Du & Wang 2011a; Du 2011a).

3. Prediction $R_{\max}$ for Cycle 24

Now, we employ the above technique to predict the peak size of cycle $n = 24$. The results are shown in Fig. 3: R_p (solid) is the predicted $R_{\max}(24)$ and r (dotted) is the correlation coefficient between $R_{\max}(i)$ and $\beta_a(\Delta m, i)$ for cycles $i = 1, 2, \dots, 23$ at a given Δm . It is seen that R_p does not vary significantly with Δm . At the current state ($\Delta m = 27$), the correlation coefficient between $R_{\max}$ and β_a is $r(27) = 0.88$, and the regression equation of $R_{\max}$ against β_a is

$$R_{\max} = 52.1 + 27.2\beta_a, \quad (4)$$

with a standard deviation of $\sigma = 19.2$. Substituting the current value of $\beta_a(27, 24) = 1.17$ into this equation, the peak sunspot number for




Fig. 3.— Prediction of $R_{\max}(24)$ as a function of Δm .

the ongoing cycle (24) is predicted as $R_p(24) = 84 \pm t_r(23)\sigma = (84 \pm 33)$ (asterisk), where $t_r(23) = 1.721$ is the t-value at the 90% level of confidence in a student's t-distribution for $n_f = 23 - 2 = 21$ degrees of freedom.

From the relationship between T_a and $R_{\max}$,

$$T_a = 79.7 - 0.251R_{\max}, \quad \sigma = 9.3, \quad (5)$$

one can estimate the rise time T_a for cycle 24. Using the predicted value (84) of $R_{\max}(24)$, one obtains $T_a(24) = (59 \pm 9)$ months. Therefore, the peak of cycle 24 may probably occur around October, 2013, slightly later than that (May, 2013) by both NASA Marshall Space Flight Center (MSFC)² based on a quasi-Planck function and NOAA space weather prediction center (SWPC)³ based on a consensus decision of “The Solar Cycle 24 Prediction Panel, and that (June, 2013) based on a modified Gaussian function (Du 2011b).

4. Discussions and Conclusions

Studying the correlation between $R_{\max}$ of a solar cycle and a related parameter is useful to understand the dynamic process of the cycle. A high correlation can be used to estimate the strength of a new solar cycle (Kane 2010; Pesnell 2008; Messerotti et al. 2009; Du 2011c; Wang et al. 2008, 2002, 2009; Le 2004; Li 2009). For example, Ohl's geomagnetic precursor method (Ohl 1976) succeeded in predicting $R_{\max}$ in cycles 20–22 (Layden 1991; Thompson 1993; Shastri 1998;

Schüssler 2007) due to the high correlation coefficients (> 0.8) between $R_{\max}$ and geomagnetic-based parameters. However, a high correlation does not always yield a satisfactory prediction (Du et al. 2009b; Du & Wang 2011a; Du 2011a; Cameron & Schüssler 2007; Du & Wang 2008, 2010) and a low correlation may also yield an accurate prediction in some cases (see Section 2.2).

A prominent feature in the solar cycle is the so-called Waldmeier effect that stronger cycles tend to rise faster (Waldmeier 1939; Hathaway et al. 2002; Du et al. 2009a; Cameron & Schüssler 2008). This effect has already begun to work in the early phase of the cycle (β_a). The variation in β_a reflects the temporal evolution of the dynamic process of solar magnetic activities from the initial phase to the peak phase of the cycle, and so, β_a can provide very useful information for the cycle.

In this study, we analyzed the temporal variation in the correlation coefficient (r) between $R_{\max}$ and β_a as a function of Δm months after the solar minimum ($R_{\min}$) and the predictive power of β_a on $R_{\max}$ as the solar cycle progresses. First, it is shown that r increases as Δm increases with an inflection point over 0.8 at about $\Delta m = 20$ months. The dynamic process of the solar activity is more non-linear near the initial phase of the cycle ($r < 0.5$ if $\Delta m \leq 10$) and tends to be stable after twenty months entering the cycle.

Besides, β_a behaved rather well in predicting $R_{\max}$: the prediction error is usually within the estimated one after about $\Delta m = 20$ months entering a solar cycle, $|R_p(\Delta m, n) - R_{\max}(n)| < t_r(n-1)\sigma(\Delta m, n-1)$ at the 90% level of confidence. This is a crucial point in prediction because a method will be less useful if the prediction is not within the prediction range derived from the method. In addition, the relative prediction error (E) based on β_a is usually less than 20% when $\Delta m \geq 20$ months. Thus, β_a is a good indicator for the subsequent $R_{\max}$. Finally, E tends to decrease as the cycle progresses. Therefore, the maximum amplitude of a new cycle ($R_{\max}$) can be well estimated at twenty months after the start.

It should be noted in Fig. 2 that the correlation between $R_{\max}$ and β_a is not strong near the initial phase of the cycle, while the prediction of $R_{\max}$ from β_a is rather good in some cycles (15, 20, 22 and 23). Therefore, a high correlation is not the sole condition to obtain a more accurate pre-

²<http://solarscience.msfc.nasa.gov/SunspotCycle.shtml>

³<http://www.swpc.noaa.gov/SolarCycle/index.html>

Table 2: Some predictions for cycle 24

Author	$R_{\max} \pm \sigma$	method or predictor
Kane (2010)	58 $\pm$ 25	<i>aa</i> minimum
Choudhuri et al. (2007)	68	solar dynamo model
NASA/MSFC	70	quasi-Planck function
Du (2011b)	72 $\pm$ 11	modified Gaussian function
Svalgaard et al. (2005)	75 $\pm$ 8	polar field
Schatten (2005)	80 $\pm$ 30	polar field
Li et al. (2005)	80 /137	slow/fast riser
Du (2011c)	82 /53	<i>aa</i> minimum/corrected
Du & Wang (2011b)	84 $\pm$ 17	similar cycles
current study	84 $\pm$ 19	rising rate
Jiang et al. (2007)	85	solar dynamo model
Li et al. (2011)	88	sunspot minimum
NOAA/SWPC	90	consensus
Wang & Sheeley (2009)	97 $\pm$ 25	open flux
Wang et al. (2009)	100 $\pm$ 8	similar cycles
Wang et al. (2002)	101 $\pm$ 18	similar cycles
Hiremath (2008)	110 $\pm$ 11	autoregression
Rigozo et al. (2011)	113	spectral components
Dabas et al. (2008)	124 $\pm$ 23	geomagnetic disturbed days
Tlatov (2009)	135 $\pm$ 12	$G \propto 1/R_z$
Du & Wang (2008)	140 $\pm$ 16	cycle length
Hathaway & Wilson (2006)	160 $\pm$ 25	I component of <i>aa</i>
Dikpati & Gilman (2006)	165 $\pm$ 15	flux-transport dynamo model

diction (Cameron & Schüssler 2007; Du & Wang 2008, 2010; Svalgaard et al. 2005; Schatten 2005), which may depend on the variation of the correlation or long-term periodicities (Du et al. 2009b; Du & Wang 2011a; Du 2011a; Du & Wang 2008, 2010). In this study, $R_{\max}$ can be well estimated from β_a even if the correlation coefficient (r) is not strong near the initial phase in some cycles, *e.g.*, $E < 20\%$ for small r (< 0.5) at small Δm in cycles 15, 20, 22 and 23 (see Fig. 2).

Based on β_a at the current state $\Delta m = 27$, the peak sunspot number of the ongoing cycle (24) is predicted to be $R_{\max} = 84 \pm 33$ at the 90% level of confidence or $R_{\max} = 84 \pm 19$ with a $1\text{-}\sigma$ uncertainty. This prediction is higher than a few predictions and lower than many others, some of which are shown in Table 2.

Accurately predicting the strength of an upcoming solar cycle is important for both solar physics and solar-terrestrial environment. A reliable prediction of $R_{\max}$ may test models for explaining the solar cycle (Pesnell 2008). So far, a great many results have been published on the prediction of $R_{\max}$ for cycle 24, of which some are based on statistics and some others are related to physics (see Table 2). As the solar activity near the minimum between cycles 23 and 24 lasts so long a time at a low level before rising (as shown in the most spotless days since cycle 16, Li et al. 2011, 2010), cycle 24 is unusual, which is drawing greater attention than ever. Besides, as solar

dynamo models have begun to be applied in predicting $R_{\max}$ (Choudhuri et al. 2007; Jiang et al. 2007; Dikpati & Gilman 2006), the predictions of the strength of cycle 24 attract special attention in order to test the predictive skill of solar dynamo models.

Discrepancies are found in the predictions of $R_{\max}$ for cycle 24 by erent methods (Table 2). Our prediction (84) is near to those by the polar field (or solar dynamo model), about 30% lower than the peak size of cycle 23. Recently, we find that cycle 24 is most likely similar to cycles 14 and 10 (Du & Wang 2011b). Therefore, even if cycle 24 is not a strong cycle, large eruption events may also occur as in cycle 10 for the largest solar storm of the year 1859 (Carrington Event).

Conclusions are summarized below.

1. The correlation coefficient (r) between the maximum amplitude ($R_{\max}$) of a solar cycle and the rising rate (β_a) at Δm months after the solar minimum ($R_{\min}$) increases as Δm increases with an inflection point at about twenty months entering the cycle.
2. The prediction error based on the linear relationship between $R_{\max}$ and β_a is usually within the estimated one when $\Delta m \geq 20$, $|R_p(\Delta m, n) - R_{\max}(n)| < t_r(n-1)\sigma(\Delta m, n-1)$, where $\sigma(\Delta m, n-1)$ is the standard deviation of the regression equation for the data up to cycle $n-1$, and $t_r(n-1)$ is the t -value at the 90% level of confidence in a student's t -distribution.
3. The relative prediction error (E) from the above technique tends to decrease as the cycle progresses and will be less than 20% when $\Delta m \geq 20$.
4. The current cycle (24) is temporarily predicted to peak around October 2013 with a size of $R_{\max} = 84 \pm 33$ at the 90% level of confidence.

Acknowledgments

The authors are grateful to the two anonymous referees for suggestive and helpful comments, which improved the original version of the manuscript. This work is supported by the National Natural Science Foundation of

China (Grant Nos. 10973020, 40890161 and 10921303), the National Basic Research Program of China (grant No. 2011CB811406), and the China Meteorological Administration (grant No. GYHY201106011).

REFERENCES

- Cameron, R., & Schüssler, M. 2007, *ApJ*, 659, 801
- Cameron, R., & Schüssler, M. 2008, *ApJ*, 685, 1291
- Choudhuri, A. R., Chatterjee, P., & Jiang, J. 2007, *Phys. Rev. Lett.*, 98, 131103
- Dabas, R. S., Sharma, K., Das, R. M., et al. 2008, *Sol. Phys.*, 250, 171
- Dikpati, M., & Gilman, P. A. 2006, *ApJ*, 649, 498
- Du, Z. L. 2011a, *Sol. Phys.*, 270, 407
- Du, Z. L. 2011b, *Sol. Phys.*, 273, 231
- Du, Z. L. 2011c, *Ap&SS*, doi: 10.1007/s10509-011-0906-4, arXiv: 1110.5973
- Du, Z. L., & Wang, H. N. 2008, *Chin. J. Astron. Astrophys.*, 8, 477
- Du, Z. L., & Wang, H. N. 2010, *RAA (Research in Astronomy and Astrophysics)*, 10, 950
- Du, Z. L., & Wang, H. N. 2011a, *Sci. Chin. Ser. G-Phys. Mech. Astron.*, 54, 172
- Du, Z. L., & Wang, H. N. 2011b, *RAA (Research in Astronomy and Astrophysics)*, 11, 1482
- Du, Z. L., Wang, H. N., Zhang, L. Y. 2009a, *Sol. Phys.*, 255, 179
- Du, Z. L., Li, R., & Wang, H. N. 2009b, *AJ*, 138, 1998
- Hathaway, D. H., & Wilson, R. M. 2006, *Geophys. Res. Lett.*, 33, L18101
- Hathaway, D. H., Wilson, R. M., & Reichmann, E. J. 2002, *Sol. Phys.*, 211, 357
- Hiremath, K. M. 2008, *Ap&SS*, 314, 45
- Jiang, J., Chatterjee, P., & Choudhuri, A. R. 2007, *MNRAS*, 381, 1527
- Kane, R. P. 2010, *Ann. Geophys.*, 28, 1463
- Layden, A. C., Fox, P. A., Howard, J. M. et al. 1991, *Sol. Phys.*, 132, 1
- Le, G. M. 2004, *Chin. J. Astron. Astrophys.*, 4, 578
- Li, K. J., Gao, P. X., & Su, T. W. 2005, *Chin. J. Astron. Astrophys.*, 5, 539
- Li, K. J. 2009, *RAA (Res. Astron. Astrophys.)*, 9, 959
- Li, K. J., Feng, W., & Liang, H. F. 2010, *Sci. Chin. Phys. Mech. Astron. (in Chinese)*, 40, 1293
- Li, K. J., Feng, W., Liang, H. F., et al. 2011, *Ann. Geophys.*, 29, 341
- Messerotti, M., Zuccarello, F., Guglielmino, S. L., et al. 2009, *Space Sci. Rev.*, 147, 121
- Ohl, A. I. 1976, *Solnice Danie*, 9, 73
- Pesnell, W. D. 2008, *Sol. Phys.*, 252, 209
- Rigozo, N. R., Souza, E. M. P., Evangelista, H. et al. 2011, *J. Atmos. Sol. Terr. Phys.*, 73, 1294
- Schatten, K. H. 2005, *Geophys. Res. Lett.*, 32, L21106
- Schüssler, M. 2007, *Astron. Nachr.*, 328, 1087
- Shastri, S. 1998, *Sol. Phys.*, 180, 499
- Svalgaard, L., Cliver, E. W., & Kamide, Y. 2005, *Geophys. Res. Lett.*, 32, L01104
- Thompson, R. J. 1993, *Sol. Phys.*, 148, 383
- Tlatov, A. G. 2009, *Ap&SS*, 323, 221
- Waldmeier, M. 1939, *Astron Mitt Zrich*, 14, 439
- Wang, Y. M. & Sheeley, N. R. 2009, *ApJ*, 694, L11
- Wang, J. L., Gong, J. C., Liu, S. Q. et al. 2002, *Chin. J. Astron. Astrophys.*, 2, 557
- Wang, J. L., Miao, J., Liu, S. Q., et al. 2008, *Sci. Chin. Ser. G-Phys. Mech. Astron.*, 51, 1938
- Wang, J. L., Zong, W. G., Le, G. M., et al. 2009, *RAA (Res. Astron. Astrophys.)*, 9, 133